

**AD-A164 241**

METAL CLUSTER TOPOLOGY 3 PLATINUM CARBONYL CLUSTERS(U)  
GEORGIA UNIV ATHENS DEPT OF CHEMISTRY R B KING  
29 JAN 86 TR-17 N00014-85-K-8365

1/1


**UNCLASSIFIED**

F/G 7/4

NL

END

<sup>a</sup>FLN is 0.



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

10

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
Technical Report #17		
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
METAL CLUSTER TOPOLOGY. 3. PLATINUM CARBONYL CLUSTERS		Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
R.B. King		N00014-85-K-0365
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
University of Georgia Department of Chemistry Athens, GA 30602		NR 051-861
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Office of Naval Research Department of the Navy Arlington, VA 22217		1/29/86
		13. NUMBER OF PAGES
		19
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
To be published in <u>Inorganica Chimica Acta</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Metal Clusters Platinum Möbius delocalization		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
<p>Previously discussed topological models of metal cluster bonding are now extended to the treatment of stacked platinum carbonyl clusters, whose structure and bonding exhibit a variety of new features. The stacked triangle cluster dianions <math>Pt_3k(CO)_6k^{2-}</math> (<math>k=2,3,4,5</math>) are best regarded as built from edge-localized bonds with additional Möbius delocalization on both the top and bottom triangles of the stack. These stacked triangles thus appear to be the first examples of stable chemical species having planar rings of atoms exhibiting Möbius rather than untwisted Hückel delocalization. Such Möbius delocalization can naturally arise from the phase changes of appropriate</p>		

DTIC  
ELECTE  
FEB 14 1988  
S D

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD-A164 241

LIB FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

#20 (Continued)

d orbitals of each of the platinum atoms in the top and bottom triangles of the stack. The more complicated tetraanion  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  can be regarded structurally as a threaded tube in which a  $\text{Pt}_{15}$  stack of three pentagons is the tube and a  $\text{Pt}_4$  chain is the thread. Edge-localized bonding is then seen to occur within both the  $\text{Pt}_{15}$  stack (25 edges) and the  $\text{Pt}_4$  thread (3 edges) with additional delocalized bonding within the pentagonal pyramidal chambers at each end of the stack. These seemingly rather exotic topological bonding models are consistent with the general principles of metal cluster bonding and reproduce exactly the observed electron counts for the stacked platinum carbonyl clusters.

S/N 0102- LR 014- 6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

OFFICE OF NAVAL RESEARCH

Contract N00014-84-K-0365

TECHNICAL REPORT NO. 17

**Metal Cluster Topology. 3. Platinum Carbonyl Clusters**

by

**R.B. King**

Prepared for Publication in

**Inorganica Chimica Acta**

University of Georgia  
Department of Chemistry  
Athens, Georgia 30602

January 29, 1986

Reproduction in whole or in part is permitted  
for any purpose of the United States Government.

This document has been approved for public release  
and sale; its distribution is unlimited.

### Abstract

Previously discussed topological models of metal cluster bonding are now extended to the treatment of stacked platinum carbonyl clusters, whose structure and bonding exhibit a variety of new features. The stacked triangle cluster dianions  $\text{Pt}_3\text{k}(\text{CO})_{6\text{k}}^{2-}$  ( $\text{k}=2,3,4,5$ ) are best regarded as built from edge-localized bonds with additional Möbius delocalization on both the top and bottom triangles of the stack. These stacked triangles thus appear to be the first examples of stable chemical species having planar rings of atoms exhibiting twisted Möbius rather than untwisted Hückel delocalization. Such Möbius delocalization can naturally arise from the phase changes of appropriate d orbitals of each of the platinum atoms in the top and bottom triangles of the stack. The more complicated tetraanion  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  can be regarded structurally as a threaded tube in which a  $\text{Pt}_{15}$  stack of three pentagons is the tube and a  $\text{Pt}_4$  chain is the thread. Edge-localized bonding is then seen to occur within both the  $\text{Pt}_{15}$  stack (25 edges) and the  $\text{Pt}_4$  thread (3 edges) with additional delocalized bonding within the pentagonal pyramidal chambers at each end of the stack. These seemingly rather exotic topological bonding models are consistent with the general principles of metal cluster bonding and reproduce exactly the observed electron counts for the stacked platinum carbonyl clusters.



Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Availability and/or Special
A-1	

## 1. Introduction

Previous papers in this series discuss the bonding topologies of osmium<sup>2</sup> and gold<sup>1</sup> clusters. This paper presents an analogous treatment of stacked platinum carbonyl clusters.<sup>3,4</sup> Such clusters are of particular interest since they have rather different structures from clusters of other transition metals. Thus whereas other transition metal clusters have structures derived from deltahedra, the stacked platinum carbonyl clusters are constructed by stacking planar rings of odd numbers of platinum atoms. Despite successes in understanding other types of metal clusters,<sup>5,6,7,8,9,10,11,12</sup> no satisfactory electron counting theory has been developed for the platinum carbonyl clusters owing at least to the following two difficulties:

(1) Uncertainties regarding the bonding orbital manifolds<sup>1</sup> of the vertex platinum atoms since cylindrical  $sp^5$  7-orbital manifolds, toroidal  $sp^2d^5$  8-orbital manifolds, and spherical  $sp^3d^5$  manifolds are all possible for platinum relating to the stabilities of 14-, 16-, and 18-electron configurations, respectively.

(2) Uncertainties regarding the number of internal orbitals<sup>9,10</sup> contributed by each vertex platinum atom to the skeletal bonding of the stacked cluster polyhedron.

These two difficulties increase the "degrees of freedom" in developing electron counting schemes for platinum clusters. However, the platinum frameworks of these clusters also have symmetry properties that restrict the allocations of electrons and orbitals thereby compensating for the above added degrees of freedom of the vertex atoms. This paper develops for the first time this symmetry approach to cluster bonding

for which the well-characterized platinum carbonyl clusters provide one of the most interesting illustrations of its applicability. An intriguing result from this work is that such symmetry considerations require stacked platinum carbonyl clusters of the type  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  to be the first examples of systems in which planar polygons of atoms exhibit Möbius rather than Hückel delocalization.



## 2. Background

The well-characterized large platinum carbonyl clusters fall into the following two categories (Figure 1):

- (1) Stacked  $\text{Pt}_3(\text{CO})_6$  triangles leading to the dianions  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  ( $k=2,3,4,5$ ).<sup>3</sup>
- (2) Three stacked  $\text{Pt}_5$  pentagons (BDB in Figure 1) having a  $\text{Pt}_4$  chain (ACCA in Figure 1) inside the stack thereby leading to the  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  cluster.<sup>4</sup>

A common feature of both of these types of systems is the stacking of  $\text{Pt}_n$  polygons leading to a system containing a principal  $C_n$  axis on which none of the platinum atoms of the polygons are located. A rotation axis of this type will be called a non-fixing axis since rotation around such an axis leaves no atoms fixed. This is the symmetry feature that will be used for the development of the bonding topology models for these systems. If the  $\text{Pt}_n$  polygons are pentagons rather than triangles, there is space for a platinum chain inside the pentagonal stack; this platinum chain is collinear with the  $C_5$  axis of the pentagonal stack.

In view of these geometrical features of the platinum carbonyl clusters, it is important to consider the properties of polyhedra of identical vertex atoms having a non-fixing  $C_n$  rotation axis. In such systems the number of electrons arising from the vertex metal atoms must be a multiple of  $n$ , or using standard mathematical notation<sup>13</sup>:

$$e_m \equiv -z(\text{mod } n) \quad (1)$$

In equation 1  $e_m$  is the number of electrons arising from the metal atom vertices and  $z$  is the charge on the species. Furthermore the number of skeletal bonding molecular orbitals in such systems divided by  $n$  must give a remainder  $r$  which corresponds to the number of skeletal bonding molecular orbitals unchanged by the  $C_n$  symmetry operation of the non-fixing axis, i.e.

$$s^+ \equiv r \pmod{n} \quad (2)$$

In equation 2  $s^+$  is the number of skeletal bonding molecular orbitals. Application of these rules to the platinum carbonyl cluster compounds allows us to make some interesting observations concerning their bonding.

Before discussing the platinum carbonyl clusters, let us illustrate these symmetry rules by their application to borane clusters which have been well-studied by other methods.<sup>5,6,14,15</sup> Table 1 illustrates these rules for the deltahedral borane dianions  $B_nH_n^{2-}$  having a non-fixing  $C_3$  axis (i.e.,  $n \equiv 0 \pmod{3}$ ). These systems are required by equation 1 to have  $2 \pmod{3}$  skeletal electrons where the 2 arises from the -2 charge on the dianion. Since each skeletal bonding orbital is occupied by an electron pair, this corresponds to a system having  $1 \pmod{3}$  skeletal bonding orbitals. This implies a unique bonding orbital located on the  $C_3$  axis. This unique bonding orbital is located at the core of the deltahedron and arises from the  $n$ -center bond of the unique internal orbitals of the vertex atoms.<sup>9,10</sup> Thus these symmetry requirements on the number of bonding orbitals in deltahedral systems having a non-fixing  $C_3$  axis are consistent with earlier metal cluster bonding models.<sup>9,10</sup>

TABLE 1

SYMMETRY RESTRICTIONS ON THE NUMBER OF SKELETAL ORBITALS  
FOR HOMOMETALLIC POLYHEDRAL CLUSTERS HAVING A NON-FIXING AXIS

Vertex Atoms	Polyhedron	Non-fixing Axis	Skeletal bonding Orbitals for a Dianion	Example
6	Octahedron	$C_3$	1 (mod 3)	$B_6H_6^{2-}$
9	Tricapped trigonal prism	$C_3$	1 (mod 3)	$B_9H_9^{2-}$
12	Icosahedron	$C_3$	1 (mod 3)	$B_{12}H_{12}^{2-}$
3k	Stacked triangles	$C_3$	1 (mod 3)	$Pt_{3k}(CO)_{6k}^{2-}$
		$C_2$	0 (mod 2)	$Pt_{3k}(CO)_{6k}^{2-}$ (k even)

### 3. Stacked Triangle Platinum Carbonyl Clusters

The rotational subgroup of the symmetry group of the stacked triangle  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters ( $k=2,3,4,5$ ) is  $D_3$  implying a  $C_3$  axis and three  $C_2$  axes in a plane perpendicular to the  $C_3$  axis. The  $C_2$  axes suggest that the skeletal molecular orbitals are the same at both the top and the bottom of the platinum triangle stack. Equation 1 indicates that the  $C_3$  axis requires  $2 \pmod{3}$  bonding electrons from the vertex atoms.

In view of these considerations a possible bonding model for the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters is a globally delocalized bonding model similar to that well-established<sup>8,9</sup> for the deltahedral  $\text{B}_n\text{H}_n^{2-}$  anions. In such a model each  $\text{Pt}(\text{CO})_2$  vertex uses three internal orbitals thereby acting as a donor of two skeletal electrons provided that the platinum atom uses a spherical  $\text{sp}^3\text{d}^5$  bonding orbital manifold.<sup>1</sup> Thus the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters are  $2n + 2$  (or  $6k + 2 \equiv 2 \pmod{6} \equiv 2 \pmod{3}$ ) skeletal electron systems like the  $\text{B}_n\text{H}_n^{2-}$  anions.

If such a bonding model is valid for the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters, then why are such clusters not deltahedra rather than stacked triangles? Overlap of the  $3k$  unique internal orbitals of the  $3k$  platinum vertices at the center of a  $\text{Pt}_{3k}$  deltahedron would give the required  $3k$ -center bond for a non-fixing  $C_3$  axis system having  $2 \pmod{3}$  skeletal bonding electrons. However, in a stacked triangle  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  globally delocalized system the unique internal orbitals of platinum atoms three or more layers apart would be too far away to overlap significantly thereby preventing the multicenter core bond required for globally delocalized bonding. Since platinum triangle stacks as high as five

layers are well characterized<sup>3</sup> (and even higher stacks may exist), a valid bonding model for these systems cannot depend upon overlap between orbitals on platinum atoms several layers apart. The bonding in these systems must therefore be localized into specific regions of the platinum network.

Such considerations make much more plausible an alternative edge-localized bonding model for the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters. A  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  system of stacked triangles has  $3 + 6(k-1) = 6k - 3$  edges which require  $12k - 6$  skeletal electrons for edge-localized bonding. Let us now consider the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters as being built from  $\text{Pt}(\text{CO})_2$  vertices using four internal orbitals noting that the distribution of carbonyl groups between terminal and bridging carbonyl groups does not affect the formal electron counting.<sup>2</sup> Such a  $\text{Pt}(\text{CO})_2$  vertex has five (i.e.,  $9 - 4$ ) external orbitals which will formally use the ten electrons from  $\text{Pt}^0$  thereby making a  $\text{Pt}(\text{CO})_2$  vertex using four internal orbitals a donor of four skeletal electrons with these electrons formally coming from the two carbonyl groups. The available skeletal electrons for the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stacked triangle clusters can then be counted as follows:

(a) $3k \text{ Pt}(\text{CO})_2$ vertices (4 internal orbitals,	
4 skeletal electrons each): $(3k)(4) =$	12k electrons
(b) -2 charge on dianion	<u>2 electrons</u>
Total skeletal electrons	12k + 2 electrons

In connection with this electron counting scheme, note the following:

(1) The vertices of interior triangles in the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stack (Figure

1) have degree four (i.e., four edges meet at each of these vertices) so that the proposed four internal orbitals from these  $\text{Pt}(\text{CO})_2$  vertices match the corresponding vertex degrees in accord with expectations for edge-localized bonding.<sup>10</sup>


(2) The vertices of the two exterior triangles in the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stack have degree three. After using three internal orbitals of these  $\text{Pt}(\text{CO})_2$  vertices for edge-localized bonding, there remains one internal orbital from each of the six platinum atoms of the two exterior triangles for further skeletal bonding. For reasons that will become apparent below, let us call this "extra" internal orbital on each vertex atom of the exterior triangles the Möbius orbital. A  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  cluster therefore has six Möbius orbitals as long as  $k$  is at least 2.

(3) Edge-localized bonding in each of the  $6k - 3$  edges of the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stack requires  $12k - 6$  skeletal electrons. Since there are a total of  $12k + 2$  skeletal electrons, this leaves eight skeletal electrons for the molecular orbitals generated from overlap of the two groups of three Möbius orbitals at the top and bottom of the triangle stack. Let us call these eight electrons the Möbius electrons. The symmetry of the  $C_2$  axes of the  $D_{3h}$   $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  systems force equal allocation of these eight Möbius electrons to the top and bottom of the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stack. This means that at both ends of the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stack we have four Möbius electrons for the molecular orbitals formed by the three triangularly situated Möbius orbitals (Figure 2). This electron counting situation suggests  $4m$  electron ( $m$  an integer) Möbius overlap involving a twisted ring of the Möbius orbitals rather than the  $4m + 2$  electron Hückel overlap found in planar aromatic hydrocarbons such as benzene.<sup>16</sup> If the Möbius orbitals are  $d$  orbitals, then this

twisted overlap is possible for an odd number of metal atoms (e.g., a triangle or pentagon but not a quadrilateral) since d orbitals change phase (i.e. "twist") at each metal center.


This bonding model for the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stacked clusters suggests edge-localized bonding along the  $6k - 3$  edges of the stack coupled with three-center delocalized Möbius bonding at both the top and the bottom of the stack. Thus the edge-localized bonding in the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters corresponds to the edge-localized carbon-carbon  $\sigma$ -bonding in benzene whereas the Möbius bonding at the top and the bottom of the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stack corresponds to the Hückel  $\pi$ -bonding in benzene.

Let us now consider the specific platinum orbitals involved in this bonding model of the  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  clusters. An individual  $\text{Pt}_3(\text{CO})_6$  triangle of the stack can be depicted as follows:



Each platinum atom forms five bonds in the plane of the triangle: two to other platinum atoms, two to bridging carbonyl groups, and one to a terminal carbonyl group. In addition to forming these five equatorial bonds the platinum atoms of the interior triangles in a  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  stack form axial bonds to the corresponding platinum atoms in the triangles above and below the interior triangle in question. These

platinum atoms are therefore seven-coordinate having approximate pentagonal bipyramidal hybridization necessarily using  $s$ ,  $x$ ,  $y$ ,  $z$ ,  $x^2-y^2$ ,  $xz$ , and  $z^2$  orbitals with the  $z$  axes taken as orthogonal to the plane of the triangle. Similarly the platinum atoms of the two exterior triangles in a  $\text{Pt}_3\text{k}(\text{CO})_6^{2-}$  stack may be regarded as six-coordinate with approximate pentagonal pyramidal hybridization, which can be taken to use  $s$ ,  $x$ ,  $y$ ,  $z$ ,  $x^2-y^2$ , and  $xy$  orbitals in order to keep the platinum hybridizations of the interior and exterior triangles as similar as possible. This leaves the  $xz$  and  $yz$  orbitals of the platinum atoms of the two exterior triangles to hybridize to generate two additional hybrids for these platinum atoms. One of these hybrids is directed to maximize in phase overlap with the Möbius orbital(s) of adjacent platinum atom(s) in the same exterior triangle. The other hybrid becomes an external orbital which contains a non-bonding electron pair already accounted for in the electron counting scheme above. The phase of the overlaps of the internal hybrids (i.e., the Möbius orbitals) in an exterior triangle can be depicted as follows:



Note the change in phase at each metal center and the single node (i.e., twist) required for Möbius rather than Hückel overlap.

This analysis shows that the platinum atoms in the  $\text{Pt}_3\text{k}(\text{CO})_6^{2-}$  have appropriate orbitals in their spherical  $sp^3d^5$  manifolds for the bonding scheme outlined above.



#### 4. The Threaded Tubular Cluster $\text{Pt}_{19}(\text{CO})_{22}^{4-}$

The structure of the threaded stacked pentagonal cluster<sup>4</sup>  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  (Figure 1) can be built as follows:

(1) Three  $\text{Pt}_5$  pentagons (BDB in Figure 1) are stacked on top of each other forming two pentagonal prismatic chambers sharing a pentagonal face. This pentagonal stack is called the  $\text{Pt}_{15}$  tube.

(2) A linear  $\text{Pt}_4$  chain (ACCA in Figure 1) is placed collinear to the  $C_5$  axis of the stacked pentagons so that the two end members of the  $\text{Pt}_4$  chain are the apices of pentagonal pyramids at the top and the bottom of the pentagonal stack and the two central members of the  $\text{Pt}_4$  chain are located in the centers of the two pentagonal prismatic chambers noted above. This  $\text{Pt}_4$  chain is called the  $\text{Pt}_4$  thread.

This structure of  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  has the following consequences:

(1) The  $\text{Pt}_4$  thread in  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  can be regarded formally as a stack of four one-vertex polygons. Using this formalism stacks of the first three types of odd-vertex polygons are found in platinum carbonyl clusters with the one- and five-vertex polygon stacks in  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  and the three-vertex polygon (i.e., triangle) stacks in the clusters  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$ .

(2) Stacks of platinum triangles and platinum pentagons may be regarded topologically as tubes. The internal volume of a tube from stacked triangles is not large enough to contain a platinum thread. However, the internal volume of a tube from stacked pentagons is large enough to contain a platinum thread as shown by the structure of  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$ . In fact the  $\text{Pt}_4$  thread inside the  $\text{Pt}_{15}$  stack of three pentagons is probably necessary to prevent collapse of the pentagonal  $\text{Pt}_{15}$  stack

into a  $Pt_{15}$  stack of five triangles. A  $Pt_{15}$  stack of three pentagons has 25 edges whereas a  $Pt_{15}$  stack of five triangles has 27 edges thereby making the latter more stable for systems which can have an appropriate number of electrons.

(3) In  $Pt_{19}(CO)_{22}^{4-}$  the center platinum atoms of the  $Pt_4$  thread located in the centers of the two pentagonal prismatic chambers have only internal orbitals. Therefore all of the electrons of these two platinum atoms are skeletal electrons. For this reason fewer carbonyl groups per platinum (i.e.,  $CO/Pt = 1.16$ ) are needed in  $Pt_{19}(CO)_{22}^{4-}$  relative to  $Pt_{3k}(CO)_{6k}^{2-}$  ( $CO/Pt=2.00$ ) in order for the system to have the necessary number of skeletal electrons.

The proposed bonding model for  $Pt_{19}(CO)_{22}^{4-}$  has the following general features:

- (1) Edge-localized bonding along the 25 edges of the  $Pt_{15}$  stack of three pentagons having the two pentagonal prismatic chambers. This requires 50 skeletal electrons.
- (2) Edge-localized bonding along the three edges of the  $Pt_4$  thread inside the pentagonal stack. This requires 6 skeletal electrons.
- (3) Delocalized bonding within the pentagonal pyramidal chambers at the top and the bottom of the pentagonal stack. These chambers can be viewed as nido systems<sup>14</sup> analogous to  $B_6H_{10}$ , which require 16 skeletal electrons ( $= 2n + 4$  for  $n = 6$ ). However, for each of the two pentagonal pyramidal chambers 10 of the required 16 skeletal electrons can come from delocalization of the five edge-localized bonds in the pentagonal face being capped to form the pentagonal pyramidal chamber. Thus the incremental number of skeletal electrons for each pentagonal pyramidal chamber is 6 ( $= 16 - 10$ ).

The 19 platinum atoms in  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  may be divided into 17 peripheral platinum atoms and two interstitial platinum atoms (C in Figure 1). All nine  $\text{sp}^3\text{d}^5$  orbitals of the spherical bonding manifold<sup>1</sup> of the interstitial platinum atoms are internal orbitals and overlap with orbitals of the platinum atoms of the  $\text{Pt}_{15}$  triple pentagonal stack by making some of the two-center edge-localized bonds in the  $\text{Pt}_{15}$  stack into three-center bonds involving an orbital of an interstitial platinum atom. The 17 peripheral platinum atoms each have four internal and five external orbitals. The internal orbitals of the peripheral platinum atoms are used as follows (see Figure 1 for the significance of the letters A, B, and D):

(A) End platinum atoms of the  $\text{Pt}_4$  thread (2 platinum atoms): three internal orbitals for a delocalized pentagonal pyramid<sup>9,10</sup> and the fourth internal orbital for a localized bond to the nearest interstitial platinum atom also in the  $\text{Pt}_4$  thread.

(B) Platinum atoms of the top and bottom  $\text{Pt}_5$  pentagons of the  $\text{Pt}_{15}$  tube (10 platinum atoms): three internal orbitals for a delocalized pentagonal pyramid<sup>9,10</sup> and the fourth internal orbital for a localized bond to the nearest platinum atom of the middle  $\text{Pt}_5$  pentagon.

(D) Platinum atoms of the middle  $\text{Pt}_5$  pentagon of the  $\text{Pt}_{15}$  tube (5 platinum atoms): all four internal orbitals are used for edge-localized bonds to neighboring platinum atoms. Note that these platinum vertices of the  $\text{Pt}_{15}$  tube have degree 4 (ignoring any interaction of the  $\text{Pt}_{15}$  tube with the  $\text{Pt}_4$  thread).

The electrons in  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  can now be counted as follows:

(1) Source of skeletal electrons:

17 PtCO vertices using 4 internal orbitals: $(17)(2)=$	34 electrons
5 "extra" CO groups	10 electrons
2 interstitial platinum atoms: $(2)(10)=$	20 electrons
-4 negative charge on anion	<u>4 electrons</u>
Total available skeletal electrons	68 electrons

(2) Use of skeletal electrons:

Edge-localized bonding in the $\text{Pt}_{15}$ tube: 25 edges	50 electrons
Edge-localized bonding in the $\text{Pt}_4$ thread: 3 edges	6 electrons
<u>Incremental</u> electrons for each of the two delocalized pentagonal pyramidal chambers: $(2)(6)=$	<u>12 electrons</u>
Total skeletal electrons required	68 electrons

This indicates that the anion  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  has exactly the number of electrons required for the above bonding model.

The anion  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  is isoelectronic with a hypothetical neutral molecule  $\text{Pt}_{19}(\text{CO})_{24}$  which would be one member (i.e.,  $n=3$ ) of a predicted series of threaded tubular clusters having the general formula  $\text{Pt}_{6n+1}(\text{CO})_{6n+6}$ . The  $\text{Pt}_{5n}$  tubes in this series of clusters consist of a stack of  $n$   $\text{Pt}_5$  pentagons. These tubes contain a thread of  $n+1$  platinum atoms. These clusters thus have  $5n+2$  peripheral platinum atoms and  $n-1$  interstitial platinum atoms. The general electron counting of the threaded tubular  $\text{Pt}_{6n+1}(\text{CO})_{6n+6}$  goes as follows:

(1) Source of skeletal electrons:

5n+2 PtCO vertices using 4 internal orbitals apiece

$$2(5n+2)=10n + 4 \text{ electrons}$$

$$n+4 \text{ "extra" CO groups} \quad 2n + 8 \text{ electrons}$$

$$n-1 \text{ interstitial platinum atoms: } 10(n-1)=\underline{10n - 10 \text{ electrons}}$$

$$\text{Total available skeletal electrons} \quad 22n + 2 \text{ electrons}$$

(2) Use of skeletal electrons:

Edge-localized bonding in the  $\text{Pt}_{5n}$  tube:

$$10n - 5 \text{ edges} \quad 20n - 10 \text{ electrons}$$

Edge-localized bonding in the  $\text{Pt}_{n+1}$  thread:

$$n \text{ edges} \quad 2n \text{ electrons}$$

Incremental electrons for each of the two

delocalized pentagonal pyramidal chambers:

$$(2)(6)=\underline{12 \text{ electrons}}$$

$$\text{Total skeletal electrons required} \quad 22n + 2 \text{ electrons}$$

Analogy with  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  suggests that these threaded tubular clusters may be found as the tetraanions  $\text{Pt}_{6n+1}(\text{CO})_{6n+4}^{4-}$  rather than as the neutral molecules  $\text{Pt}_{6n+1}(\text{CO})_{6n+6}$ .

5. Summary

This paper presents the first detailed electron-precise bonding models of the previously enigmatic platinum carbonyl clusters  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  ( $k \geq 2$ ) and  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$ . These systems are seen to consist of stacks of platinum atoms, platinum triangles, and/or platinum pentagons having edge-localized bonding within the stacks. The tops

and bottoms of the platinum stacks in these clusters contain interesting additional features, namely Möbius triangles in the cases of the triangular stacks  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  and delocalized pentagonal pyramids in the case of the pentagonal stack  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$ . The  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  systems are significant in representing the first series of stable chemical species in which orbitals from a planar group of atoms appear to overlap in a twisted Möbius manner rather than in an untwisted Hückel manner; symmetry considerations require this unusual model made feasible by the properties of appropriate platinum d orbitals. The  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$  cluster appears to be the first known example of a threaded tubular cluster derived from a neutral species  $\text{Pt}_{6n+1}(\text{CO})_{6n+6}$  having a tube consisting of a  $\text{Pt}_{5n}$  pentagonal stack threaded by a  $\text{Pt}_{n+1}$  chain. The synthesis of other threaded tubular clusters of this series represents an interesting challenge.

Acknowledgment: I am indebted to the Office of Naval Research for partial support of this research.


## REFERENCES

- (1) For part 2 of this series see King, R.B. submitted for publication.
- (2) King, R.B. submitted for publication.
- (3) Calabrese, J.C.; Dahl, L.F.; Chini, P.; Longoni, G.; Martinengo, S. J. Am. Chem. Soc. **1974**, 96, 2614.
- (4) Washecheck, D.M.; Wucherer, E.J.; Dahl, L.F.; Ceriotti, A.; Longoni, G.; Manessero, M.; Sansoni, M.; Chini, P. J. Am. Chem. Soc. **1979**, 101, 6110.
- (5) Wade, K. Chem. Comm. **1971**, 792.
- (6) Grimes, R.N. Ann. N.Y. Acad. Sci. **1974**, 239, 180.
- (7) Wade, K. Adv. Inorg. Chem. Radiochem. **1976**, 18, 1.
- (8) Stone, A.J. Inorg. Chem. **1981**, 20, 563.
- (9) King, R.B.; Rouvray, D.H. J. Am. Chem. Soc. **1977**, 99, 7834.
- (10) King, R.B. in "Chemical Applications of Topology and Graph Theory," King, R.B., Ed; Elsevier: Amsterdam, 1983, pp. 99-123.
- (11) Teo, B.K. Inorg. Chem. **1984**, 23, 1251.
- (12) Teo, B.K.; Longoni, G.; Chung, F.R.K. Inorg. Chem. **1984**, 23, 1257.
- (13) Barnett, I.A. "Elements of Number Theory," Prindle, Weber, and Schmidt: Boston, Mass., 1972, pp. 79-81.
- (14) Rudolph, R.; Pretzer, W.R. Inorg. Chem. **1972**, 11, 1974.
- (15) Williams, R.E. Adv. Inorg. Chem. Radiochem. **1976**, 18, 67.
- (16) Zimmerman, H.E. Accts. Chem. Res. **1971**, 4, 272.


Figure 1: Schematic diagrams of the stacked triangle platinum carbonyl anion clusters  $\text{Pt}_{3k}(\text{CO})_{6k}^{2-}$  and the threaded tubular cluster  $\text{Pt}_{19}(\text{CO})_{22}^{4-}$ .

Figure 2: A comparison of the energy levels in Hückel and Möbius triangles.








$$k=2$$




$$k=3$$





**Hückel Overlap**  
(no phase change)



**Möbius Overlap**  
(odd number of phase changes)

**END**

**FILMED**

**386**

**DTIC**